

UNIVERSITI TEKNOLOGI MARA

**THE INFLUENCE OF GENETIC
POLYMORPHISMS OF CYP3A ON
THE PHARMACOKINETIC
VARIATION AND CLINICAL
OUTCOMES OF TACROLIMUS IN A
COHORT OF PATIENTS**

MOHD RAHIMI BIN MUDA

Thesis submitted in fulfillment
of the requirements for the degree of
Master of Science

Faculty of Pharmacy

February 2014

AUTHOR'S DECLARATION

I declare that the work in this thesis was carried out in accordance with the regulations of University Teknologi MARA. It is original and the result of my own work, unless otherwise indicated or acknowledged as referenced work. This thesis has not been submitted to any other academic or non-academic institution for any other degree or qualification.

I, hereby acknowledge that I have been supplied with the Academic Rules and Regulations for Postgraduate, Universiti Teknologi MARA, regulating the conduct of my study and research.

Name of Student : Mohd Rahimi Bin Muda

Student's ID No : 2009763155

Programme : Master of Science (PH780)

Faculty : Pharmacy

Title : The Influence of Genetic Polymorphism of *CYP3A* on The Pharmacokinetic Variation and Clinical Outcomes of Tacrolimus in a Cohort of Patients

Signature of Student :.....

Date : February 2014

ABSTRACT

The complexity of determining the variability in tacrolimus pharmacokinetics parameter among organ transplant patients and its narrow therapeutic window characteristics necessitates proper drug monitoring in order to ensure the level is maintained within therapeutic range. Many studies have reported that variability of tacrolimus pharmacokinetics was due to differential expression of drug metabolizing enzymes CYP3A4 and CYP3A5 encoded by *CYP3A4* and *CYP3A5* genes respectively. The aim of the study was to develop a genotyping method in detecting *CYP3A4*18*, *CYP3A5*3* and *CYP3A5*6* mutations as well as investigating any correlation of genotyping findings with the clinical and tacrolimus related pharmacokinetics data and to quantify enzyme expression through the relationship of *CYP3A4* and *CYP3A5* mRNA copy number with the patients' genotypes. Five ml of blood were withdrawn from the 80 patients and the DNA was extracted, thus proceeded for genotyping of *CYP3A4*18*, *CYP3A5*3* and *CYP3A5*6* variants using two steps PCR technique. Absolute quantification was employed where the serial dilution of plasmid from cloned bacteria to generate a standard curve was done prior to real time PCR method. Nine of 80 patients were found to be heterozygous of *CYP3A4*18* (11.25%). As result of *CYP3A5*3* polymorphism, eight patients were found as homozygous wild type (10%), 31 patients were heterozygous (38.75%) and 41 patients were homozygous mutant (51.25%). No *CYP3A5*6* mutation was observed in this study. Homozygous mutant was significantly associated with lowest requirement of tacrolimus dose and highest adjusted level per dose ($P < 0.01$). There was no significant correlation between *CYP3A4* and *CYP3A5* mRNA copy number with the *CYP3A4*18* and *CYP3A5*3* genotypes respectively. Significant correlation between *CYP3A5*3* genotypes with the doses has provided a reliable information in predicting the appropriate individual tacrolimus doses.

ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful

All praise to Allah for His limitless and continuous blessing in completing this thesis. First and foremost, I would like to thank to my mentor and supervisor, Professor Dr Teh Lay Kek for her priceless guidance, patience, advice and supervision in pursuing this project. Not to forget, my appreciation also goes to my co-supervisor, Professor Dr Mohd Zaki Salleh for all his constructive comments, knowledge and idea relating to the topic of this study.

I would like to express my appreciation to the Ministry of Science, Technology and Innovation Malaysia for granting the research grant (Grant number: 02-01-01-SF0171). My thankfulness also goes to UiTM and Professor Dr Aishah Adam for awarding the scholarship as well as giving the opportunity to pursue my study in the Faculty of Pharmacy. Extended appreciation goes to all of the research collaborators and staffs from Nephrology and Urology department of Hospital Kala Lumpur and Hospital Selayang for their assistance during the sample collection.

Sincere heartfelt thanks goes to all member of Pharmacogenomic Centre (PROMISE), UiTM especially Ikhwan, Siti Nasirah, Lee Lian Shien, Fazleen Haslinda, Dr Sharina and Nor Izwani for their moral support and kindness within the study period. Lastly, my deepest gratitude and love to my family for their unconditional love, support and prayers throughout my life. Thank you!.

TABLE OF CONTENTS

AUTHOR'S DECLARATION	ii
ABSTRACT	iii
ACKNOWLEDGEMENT	iv
TABLE OF CONTENTS	v
LIST OF TABLES	xi
LIST OF FIGURES	xv
LIST OF PLATES	xviii
LIST OF EQUATIONS	xix
LIST OF ABBREVIATIONS	xx

CHAPTER ONE: INTRODUCTION

1.1	Background of the study	1
1.2	Statement of research problems	2
1.3	Objectives	3

CHAPTER TWO: LITERATURE REVIEW

2.1	History of transplantation and usage of immunosuppression therapy	4
2.2	Tacrolimus	5
2.2.1	Structure and physicochemical properties of tacrolimus	6
2.3	Pharmacodynamic of tacrolimus	7
2.4	Pharmacokinetics of tacrolimus	8
2.4.1	Absorption and Distribution	8
2.4.2	Metabolism and Excretion	8
2.5	Therapeutic Drug Monitoring of Tacrolimus	9
2.5.1	Overview	9
2.5.2	Determination of initial dosage method	10
2.5.3	Limitation in therapeutic drug monitoring	12